MEFTAL-SPAS











Dicycloverine / Dicyclomine(10.0 Mg)

Mefenamic Acid(250.0 Mg)

USES

- ☐ Menstrual period cramps
- ☐ Irritable bowel syndrome
- Heavy bleeding during periods
- ☐ Muscle pain and inflammation
- ☐ Fever and joint pain
- ☐ Stomach and abdominal pain

Dosage

>For adults

Initially, 500 mg is given once, followed by 250 mg every 6 to 8 hours. The scheduled dose should not exceed more than three days.

>For children≥ 14 years:

Initially, 500 mg is given once, followed by 250 mg every 6 hourly. The scheduled dose should not exceed more than 7 days.

- ➤ Meftal Spas tablets should be taken as directed by your doctor.
- > Do not cut, break or chew the medicine.
- > It can be taken with or after food.
- ➤ It would be best if you took it at a fixed time for optimal results.
- ➤ Do not consume it more than that prescribed by your doctor.

- Can I take Meftal-Spas tablet during pregnancy?
- Meftal-Spas tablet should not be used in a pregnant woman due to lack of safety information, especially in last trimester of pregnancy.

- Can I take Meftal-Spas tablet while breastfeeding?
- Breast-feeding must be discontinued during treatment with Meftal-Spas tablet.

- Is it safe to prescribe this medicine to people with high or low blood pressure?
- Consuming this medicine may cause fluid retention and high blood pressure.
- Therefore, patients with hypertension or even hypotension should refrain from using this medicine without a doctor's consultation.

Side effects

- Nausea
- **Vomiting**
- Stomach upset
- **Dizziness**
- **Drowsiness**
- Dry mouth
- Blurred vision
- Weakness

