

JAPC/18

Sl. No. :



Register
Number

--	--	--	--	--	--	--	--	--	--

2018

PHARMACEUTICAL CHEMISTRY
(Degree Standard)

Time Allowed : 3 Hours]

[Maximum Marks : 300


Read the following instructions carefully before you begin to answer the questions.

IMPORTANT INSTRUCTIONS

1. The applicant will be supplied with Question Booklet 15 minutes before commencement of the examination.
2. This Question Booklet contains 200 questions. Prior to attempting to answer the candidates are requested to check whether all the questions are there in series and ensure there are no blank pages in the question booklet. **In case any defect in the Question Paper is noticed it shall be reported to the Invigilator within first 10 minutes and get it replaced with a complete Question Booklet. If any defect is noticed in the Question Booklet after the commencement of examination it will not be replaced.**
3. Answer **all** questions. All questions carry equal marks.
4. You must write your Register Number in the space provided on the top right side of this page. Do not write anything else on the Question Booklet.
5. An answer sheet will be supplied to you, separately by the Room Invigilator to mark the answers.
6. You will also encode your Question Booklet Number with Blue or Black ink Ball point pen in the space provided on the side 2 of the Answer Sheet. If you do not encode properly or fail to encode the above information, action will be taken as per commission's notification.
7. Each question comprises *four* responses (A), (B), (C) and (D). You are to select **ONLY ONE** correct response and mark in your Answer Sheet. In case you feel that there are more than one correct response, mark the response which you consider the best. In any case, choose **ONLY ONE** response for each question. Your total marks will depend on the number of correct responses marked by you in the Answer Sheet.
8. In the Answer Sheet there are **four** circles (A), (B), (C) and (D) against each question. To answer the questions you are to mark with Blue or Black ink Ball point pen **ONLY ONE** circle of your choice for each question. Select one response for each question in the Question Booklet and mark in the Answer Sheet. If you mark more than one answer for one question, the answer will be treated as wrong. *e.g.* If for any item, (B) is the correct answer, you have to mark as follows :
(A) ● (C) (D)
9. You should not remove or tear off any sheet from this Question Booklet. You are not allowed to take this Question Booklet and the Answer Sheet out of the Examination Hall during the time of examination. After the examination is concluded, you must hand over your Answer Sheet to the Invigilator. You are allowed to take the Question Booklet with you only after the Examination is over.
10. The sheet before the last page of the Question Booklet can be used for Rough Work.
11. Do not tick-mark or mark the answers in the Question Booklet.
12. Applicants have to write and shade the total number of answer fields left blank on the boxes provided at side 2 of OMR Answer Sheet. An extra time of 5 minutes will be given to specify the number of answer fields left blank.
13. Failure to comply with any of the above instructions will render you liable to such action or penalty as the Commission may decide at their discretion.

SEAL

SPACE FOR ROUGH WORK



PHARMACY --- INDIA

1. Silver nitrate is assayed by
 (A) Precipitation method
 (B) Complexometric method
 (C) Acidimetry
 (D) Gravimetry method
2. _____ is prepared by dissolving aluminium hydroxide in sulphuric acid.
 (A) Aluminium sulphate
 (B) Aluminium chloride
 (C) Aluminium oxide
 (D) Aluminium sulphite
3. _____ is prepared by neutralization of sodium carbonate with hydrogen fluoride.
 (A) Sodium formamide
 (B) Sodium fluoride
 (C) Sodium bicarbonate
 (D) Hydro fluoric acid
4. The number of protons in an elements is equal to its
 (A) Atomic number
 (B) Neutron number
 (C) Ionic number
 (D) Mass number
5. The nuclides that have same atomic number and different mass numbers are called
 (A) isobars
 (B) isotopes
 (C) isotones
 (D) nuclides
6. _____ is added to make alkaline which in turn stabilizes the complex in the limit test for Iron.
 (A) NaOH
 (B) Ammonia
 (C) NaHCO₃
 (D) Ammonium chloride

7. _____ is used in the preparation of poultices.
- (A) Bentonite (B) Heavy Kaolin
(C) Charcoal (D) Light Kaolin
8. In complexometric titration, to set a sharp endpoint, a known amount of _____ is added.
- (A) Magnesium oxide (B) Magnesium carbonate
(C) Magnesium sulphate (D) Magnesium sulphite
9. _____ is used for sterilizing surgical instruments.
- (A) Cesium - 137 (B) Cobalt - 60
(C) Calcium - 110 (D) Mercury - 203
10. _____ theory is used to explain the nature of bonding in coordination compounds.
- (A) Lewis Theory (B) Gibb's Theory
(C) Werner's Theory (D) Hess Theory
11. The phenomenon of maintenance of body fluids and the concentration of different electrolytes in extracellular and intracellular fluids is known as
- (A) Osmotic equilibrium
(B) Blood volume
(C) Homeostasis
(D) Acid-base balance
12. Which of the following acids is used for the preparation of ceric ammonium sulphate I.P?
- (A) Hydrochloric acid
(B) Sulphuric acid
(C) Nitric acid
(D) Perchloric acid

13. Ammonium carbonate is used as
(A) Antacid
(C) Antidote
(B) Respiratory stimulant
(D) Hematinic
14. Calcium folinate is assayed by
(A) Spectro photometry
(C) HPTLC
(B) HPLC
(D) Spectro fluorimetry
15. Aluminium chloride is assayed by
(A) Precipitation titration
(B) Non-aqueous titration
(C) Gravimetric method
(D) Complexometric titration
16. Potassium in oral rehydration salts is determined by
(A) Polarimetry
(C) HPLC
(B) UV-Spectrometry
(D) Flame photometry
17. Sodium citrate is assayed by
(A) Precipitation titration
(C) Complexometric method
(B) Gravimetry method
(D) Non-aqueous titration
18. Ferrous succinate is assayed by
(A) Redox titration
(C) Non-aqueous titration
(B) Precipitation titration
(D) Gravimetry method
19. Penicillamine is assayed by
(A) Precipitation titration
(C) Gravimetry method
(B) Non-aqueous titration
(D) Redox titration

20. Which one of the following statements is not true about iodine value in the analysis of oils?
- (A) It can be used to detect adulteration in oils
(B) It is directly proportional to the drying power of the oil
(C) It is the measure of saturated acid present in the oil
(D) It is the measure of unsaturated acid present in the oil
21. The distinction between waxes and fats is based entirely on
- (A) Chemical composition (B) Melting point
(C) Density (D) Refractive index
22. Waxes are similar to fats, but they are more difficult to
- (A) Solidity (B) Saponify
(C) Dissolve in non-polar solvents (D) Melt by heat
23. If sesame oil is adulterated, it is tested by the _____ reaction.
- (A) Baudouin (B) Kolbe's
(C) Reimer-tiemann (D) Sand meyer
24. Iodine value gives an indication of the proportion of _____ in the substance.
- (A) Mineral acid (B) Carboxylic acid
(C) Saturated acid (D) Unsaturated acid
25. Acid value is the number of milligram of Potassium hydroxide required to neutralise the free acid in 1 g. of the
- (A) Carbohydrate (B) Proteins
(C) Vitamins (D) Fats
26. In the determination of magnesium sulphate by complexometric method, which of the following buffer is used?
- (A) Acetate buffer pH 4.0 (B) Acetate buffer pH 5.0
(C) Ammonia buffer pH 10.0 (D) Phosphate buffer pH 8.0

27. The molecular weight "M" of the solute can be calculated from osmotic pressure by using the formula

(A) $M = wRT / \pi V^2$

(B) $M = wRT^2 / \pi V$

(C) $M = wRT / \pi V$

(D) $M = wRT / \pi^2 V^2$

28. Match the following :

- | | |
|-------------------------|--|
| (a) Isotonic solution | (i) Lower osmotic pressure than the other |
| (b) Hypertonic solution | (ii) Different osmotic pressure |
| (c) Hypotonic solution | (iii) Higher osmotic pressure than the other |
| | (iv) Same osmotic pressure |

- | | | |
|--|-------|-------|
| (a) | (b) | (c) |
| (A) (ii) | (iv) | (iii) |
| <input checked="" type="checkbox"/> (B) (iv) | (iii) | (i) |
| (C) (iii) | (i) | (iv) |
| (D) (i) | (iii) | (ii) |

29. The preparation of an optical active chiral compound from Achiral molecule under the influence of optical active substance is known as

- | | |
|-----------------------|--|
| (A) Achiral synthesis | <input checked="" type="checkbox"/> (B) Asymmetric synthesis |
| (C) Retro synthesis | (D) Combinatorial synthesis |

30. Sodium lactate is assayed by

- | | |
|-----------------------------|---|
| (A) Non-aqueous titration | <input checked="" type="checkbox"/> (B) Acid base titration |
| (C) Precipitation titration | (D) Gravimetric method |

31. When one of the products of a reaction itself act as a catalyst for that reaction the phenomenon is called as

- | | |
|-----------------------|--|
| (A) Poison catalyst | <input checked="" type="checkbox"/> (B) Auto catalysis |
| (C) Negative catalyst | (D) Positive catalyst |

32. Absolute configuration is based on
 (A) Cahn-Ingold-Prelog Rule (B) Van't hoff theory
(C) Label's theory (D) Both (B) and (C)
33. How many stereo isomers are possible for a compound which has three chiral carbon atom?
(A) 6 (B) 8
(C) 10 (D) 12
34. The ratio of the velocity of light in vaccum or air to that in the substance is called as
(A) Optical rotation (B) Refractive index
(C) Polarisation (D) Viscosity
35. Liquid mixture which distil with a change in composition are called as
(A) Phase rule
(B) Azerotropic solution
 (C) Zerotropic mixture
(D) Fractional distillation
36. A substance which alters the rate of a chemical reaction and itself remaining chemically unchanged at the end of the reaction is called as
 (A) Catalyst (B) Phase rule
(C) Osmatic pressure (D) Polarisation
37. The change in enthalpy when one mole of a substance is dissolved in a specified quantity of solvent at a given temperature is defined as
(A) Heat of formation
 (B) Heat of solution
(C) Heat of combustion
(D) Heat of neutralisation

38. Vitamin E prevents rancidity by virtue of its _____ property.

- (A) Anti oxidant (B) Oxidant
(C) Sulfuration (D) Hydrogenation

39. Which of the following is a fat soluble vitamin?

- (A) Folic acid (B) B
(C) C (D) A

40. Which one of the following is temporary effect in the presence of an attacking reagent?

- (A) Mesomeric effect (B) Inductive effect
 (C) Electromeric effect (D) Isomeric effect

41. Which of the following order of Electronegativity is correct?

- (A) $O > N > Cl > F > Br$
(B) $Br > F > Cl > N > O$
(C) $C, H > Br > Cl, N > O > F$
 (D) $F > O > Cl, N > Br > C, H$

42. The C — H bond length decreases in the order given below

- (A) $C_{Sp^3} - H > C_{Sp^2} - H > C_{Sp} - H$
(B) $C_{Sp} - H > C_{Sp^2} - H > C_{Sp^3} - H$
(C) $C_{Sp^2} - H > C_{Sp^3} - H > C_{Sp} - H$
(D) $C_{Sp^3} - H < C_{Sp^2} - H < C_{Sp} - H$

43. Which of the following is Aceto acetic ester?

- (A) $CH_3 CO CH_2 COO C_2H_5$
(B) $C_2H_5 COO CH_2 COO C_2H_5$
(C) $C_2H_5 CO CH_2 COO C_2H_5$
(D) $C_2H_5 CO CH_2 CH_2 COO C_2H_5$

44. Cardiac glycosides are composed of
- (A) Sugar portion only
 - (B) Non sugar portion only
 - (C) -COOH and -OH
 - (D) Glycone and Aglycone
45. _____ is responsible for cardiotoxic activity of cardiac glycosides.
- (A) Glycone
 - (B) Aglycone
 - (C) Terpenes
 - (D) Flavones
46. In cyclopentano perhydro phenanthrene ring of cardiac glycosides ring A/B and C/D are
- (A) Cis - fused
 - (B) Trans - fused
 - (C) Cis - trans fused
 - (D) R - S fused
47. In ethylene $C = C$, one bond is _____ and other bond is _____.
- (A) Pi and Pi
 - (B) Sigma and Sigma
 - (C) Sigma and Pi
 - (D) Delta and Delta
48. Whenever carbon is bonded to four other atoms or groups it is through
- (A) Sp^2 hybrid orbitals
 - (B) Sp hybrid orbitals
 - (C) Sp^3 hybrid orbitals
 - (D) Sp^4 hybrid orbitals
49. The bonds formed by transfer of Valence electrons from one atom to another is called
- (A) Co-ordinate bonds
 - (B) Co-valent bonds
 - (C) n bonds
 - (D) Ionic bond (or) electrovalent bond


50. _____ is the spiroimidazo - piperidyl derivative of rifamycin B.

- (A) Ethionamide (B) Capreomycin
(C) Rifabutin (D) Cycloserine

51. Which one of the following compound is first generation quinolone derivative?

- (A) Isoniazid (B) Chloroquine
(C) Chloramphenicol (D) Nalidixic acid

52. Fill in the Elimination reaction.



- (A) $\text{CH}_3 - \text{OH}$
(B) $\text{CH}_3 - \text{CH}_2 - \text{COOH}$
(C) $\text{CH}_2 = \text{CH}_2$
(D) CH_4

53. The basis of antibacterial action of β -lactams is that these drugs become bound to what portion of the cell wall?

- (A) Penta glycine (B) Transpeptidase (PBP)
(C) Mycolic acid (D) D - alanine - D - alanine

54. Which one of the following is classified as antifungal antibiotics?

- (A) Flucytosine (B) Sordarin
(C) Amphotericin B (D) Resorcinol

55. The echino candins are effective systemic antifungal agents by virtue of their ability to

- (A) Block cell membrane synthesis at ergosterol level.
(B) Damage the cell membrane leading to 'leakage' of membrane
(C) Block cell membrane synthesis at β - glucan level
(D) Interfere with zymosterol synthesis

56. _____ is a measure of how readily a compound is able to attack an electron deficient atom.
- (A) Electrophilicity
(B) Nucleophilicity
(C) Basicity
(D) Acidity
57. An electrophile is
- (A) Positive charged species
(B) Negative charged species
(C) Neutral charged species
(D) Have zero charge
58. The aromaticity of an organic compound is governed by
- (A) Martownikoff's rule
(B) Crum Brown and Gibson's rule
(C) Huckel's $4n + 2$ rule
(D) Hund's rule
59. Sulphonamide contain _____ functional group
- (A) $\text{NH}_2 - \text{CO} - \text{NH}_2$
(B) $-\text{SO}_3\text{H}$
(C) $-\text{SO}_2\text{NH}_2$
(D) $\text{N} \equiv \text{N}$
60. Molecule is called as laughing gas
- (A) Nitric oxide
(B) Nitrous oxide
(C) Anaesthetic ether
(D) Cyclo propane
61. Phenobarbital is _____ acting barbiturates
- (A) Ultra short
(B) Short
(C) Intermediate
(D) Long

62. In chloroquine C₃ or C₈ alkyl group _____ the action.
(A) Increases (B) Decreases
(C) No change (D) Loss of action
63. Identify the one major opioid receptor given below
(A) α - alpha (B) β - Beta
(C) γ - gamma (D) δ - Delta
64. Phenobarbital requires dose adjustments when co-administered with
(A) Pentobarbital (B) Flurazepam
 (C) Phenytoin (D) Estazolam
65. Which one of the following is an Intravenous general anesthetics
(A) Halothane (B) Isoflurane
(C) Desflurane (D) Ketamine
66. Codeine is produces _____ pharmacological effect
(A) Anesthetic
(B) Antipyretic
(C) Antibacterial
 (D) Analgesic and Antitussive
67. (+) Isoflurane is _____ more potent than (-) Isoflurane
(A) 25% (B) 50%
(C) 75% (D) 100%
68. Which one of the following is a Morphine analogue?
(A) Levorphanol
(B) Dextro methorphan
 (C) Codeine
(D) Butorphanol

69. _____ is an optically active compound which occur in wool fat along with its derivative dihydrolanosterol
- (A) Ergosterol
(B) Lanosterol
(C) Stigmasterol
(D) Cholesterol
70. Which of the following steroids has an aromatic Ring A?
- (A) Testosterone
(B) Oestrone
(C) Progesterone
(D) Aldosterone
71. The bile acids are first converted into _____ acid which are degraded upto aetiobiliaric acid.
- (A) Isolithobiliaric
(B) Choleric
(C) Lithobiliaric
(D) Lithocholic
72. _____ may be defined as 'the critical and typical monomer units of the peptides and proteins
- (A) Oxytocin
(B) Amino acids
(C) Carbohydration
(D) Lipid
73. Which one of the following possesses Antimalarial activity?
- (A) Ephedrine
(B) Caffeine
(C) Berberine
(D) Cinchonine
74. _____ method helps to identify and determine the number of methyl moiety directly attached to N-atom.
- (A) Kuhn-Roth method
(B) Hoffmann exhaustive method
(C) Herzig method
(D) Gibbs method

75. The starting material for Strecker's synthesis of α - amino acids is
- (A) A Ketone
(B) Carboxylic acid
(C) Ethyl malonate
(D) An alkyl aldehyde
76. An example of keto-hexoses is
- (A) Lactose
(B) Fructose
(C) Glucose
(D) Mannose
77. Sucrose on treatment with the enzyme invertase gives equimolar mixture of
- (A) D(+) Galactose & D(-) Fructose
(B) D(-) Glucose & D(-) Fructose
(C) D Mannose & D(-) Fructose
(D) D Glucose & D Galactose
78. Atropine is the ester of
- (A) Tropine and Mandelic acid
(B) Tropine and Tropic acid
(C) Tropic acid and Salicylic acid
(D) Tropine and Salicylic acid
79. The double bond present in progesterone is in the _____ position.
- (A) 18 - 19
(B) 5 - 6
(C) 4 - 5
(D) 11 - 12
80. Polymers like agarose and polyacrylamides, in which the macromolecules are cross linked to give a porous _____ structure.
- (A) One dimensional
(B) Two dimensional
(C) Three - dimensional
(D) Four dimensional

81. The most commonly used cells in UV are made of
- (A) Glass
 - (B) Quartz or fused silica
 - (C) Graphite
 - (D) Sodium chloride
82. The UV – Visible region of electro magnetic lies between
- (A) 200 nm – 800 nm
 - (B) 800 nm – 1000 nm
 - (C) 1 nm – 100 nm
 - (D) 100 nm – 200 nm
83. Highly fluorescent substances have quantum yield of fluorescence (ϕ) values near
- | | |
|---|---------|
| (A) 0 | (B) 0.5 |
| <input checked="" type="checkbox"/> (C) 1 | (D) 0.2 |
84. In spectrofluorometric estimation, wide slit (width) setting introduce problem due to
- (A) Thermal decomposition
 - (B) Chemical decomposition
 - (C) Photochemical decomposition
 - (D) Physical decomposition
85. Quenching means
- (A) Reduction of viscosity
 - (B) Reduction of Adsorption
 - (C) Reduction of excitation
 - (D) Reduction of fluorescence intensity

86. _____ is the time it takes for an unretained species to pass through a chromatographic column.
- (A) Retention time
(B) Void time
(C) Flow time
(D) Run time
87. The grades of silica gel 60 (or) 140 (or) 150 indicates
- (A) Particle size
(B) Mean pore size
(C) Particle shape
(D) Price based on grades
88. HPLC columns are made up of
- (A) Glass
(B) Stainless steel (highly polished surface)
(C) High quality fiber
(D) Aluminium
89. Which one of the following detector is employed in HPLC?
- (A) Flame ionisation detector
(B) Electron capture detector
(C) Refractive index detector
(D) Thermal conductivity detector

90. The vibrational frequencies in infrared spectrophotometry is calculated by
- (A) Beer law
 - (B) Hooke's law
 - (C) Absorption law
 - (D) Lambert's law
91. The most commonly used mulling agent is
- (A) Crystalline sodium chloride
 - (B) Sodium chloride
 - (C) Nujol
 - (D) Primary amines
92. Tetra methyl silane is employed as a reference compound for
- (A) IR
 - (B) NMR
 - (C) Mass
 - (D) UV
93. In NMR spectroscopy, the radiation used for nuclear excitation is
- (A) Radio wave
 - (B) Micro wave
 - (C) UV
 - (D) IR

94. Measurement of conductance obtained in the region of the equivalence point are due to _____ of the reaction product.
- (A) Oxidation
(B) Reduction
(C) Hydrolysis
(D) Association
95. _____ is a separation method based on the differential rates of migration of charged species in an applied dc electric field.
- (A) Ion exchange
(B) Gel filtration
(C) HPLC
(D) Electrophoresis
96. For a precipitation titration, the following technique is best for detecting end point
- (A) Amperometry
(B) Diaro method
(C) Conductometry
(D) Solochrome black indicator
97. The reciprocal of resistance $\frac{1}{R}$ for an electrolytic solution is
- (A) Electrolyte solution
(B) Ion selectivity
(C) Liquid liquid electrodes
(D) Conductance

98. Potassium also has an isotope _____ which has a long life and occurs naturally mixed with stable isotope ^{39}K .
- (A) ^{38}K
- (B) ^{40}K
- (C) ^{37}K
- (D) ^{36}K
99. Gold (^{198}Au) injection is used for
- (A) Diagnosis of RBC count
- (B) Diagnosis of Blood circulation in liver
- (C) Diagnosis of Thyroid function
- (D) Treatment of Thyroid disorder
100. Which one of the following compounds is used for diagnosis of renal function
- (A) Sodium rose Bengal injection
- (B) Sodium phosphate injection
- (C) Sodium Iodide injection
- (D) Sodium Iodotripropate ^{131}I injection
101. The time required for one half of a given number of atoms to decay is called
- (A) Half life
- (B) Radio activity
- (C) AUC
- (D) Elimination rate constant
102. Titanium dioxide is assayed by
- (A) Precipitation titration
- (B) Acidimetry
- (C) Complexometric titration
- (D) Gravimetry method

103. Zinc undecylenate is assayed by
 (A) Acidimetry – Alkalimetry method
(B) Complexometric method
(C) Non aqueous titration
(D) Precipitation titration
104. _____ is used to dissolve other impurity in the limit test for chloride.
 (A) Dilute Nitric acid (B) Dilute Sulphuric acid
(C) Dilute Acetic acid (D) Dilute Silver nitrate
105. Zinc chloride is assayed by
 (A) Complexometry titration (B) Alkalimetry
(C) Acidimetry (D) Non-aqueous titration
106. Radio activity can be detected by the use of
 (A) Flame photometer
(B) Geiger Muller counter
(C) Nephelo-Turbidometer
(D) Spectrophotometer
107. Sodium meta phosphate is used as
(A) Desensitizing agent (B) Antacid
 (C) Dentifrice (D) Antidote
108. Strontium chloride is used in treatment of
(A) Conjunctivities (B) Gastric acidity
(C) Constipation (D) Dental hypersensitivity

109. _____ can be obtained by neutralization of hydrochloric acid with lime.
- (A) Calcium carbonate
 (C) Calcium chloride
(B) Calcium oxide
(D) Calcium gluconate
110. _____ is prepared by sublimation of ammonium chloride with calcium carbonate.
- (A) Ammonium oxalate
 (B) Ammonium carbonate
(C) Alum
(D) Aluminium hydroxide
111. Which of the following drug is used as diuretic?
- (A) Sodium citrate
 (C) Ammonium chloride
(B) Potassium citrate
(D) Sodium acetate
112. Disodium Edetate is used as
- (A) Expectorant
(C) Respiratory stimulant
 (D) Anti dote for metal Poisoning
(B) Emetic
113. Dimercaprol is used as
- (A) Antidote for heavy metal poisoning
(C) Acidifier
(B) Antacid
(D) Laxative
114. Ferrous gluconate is used as
- (A) Antacid
(C) Systemic acidifier
 (B) Haematinic agent
(D) Antidote
115. Which one of the following is used in the treatment of Wilson's disease?
- (A) Dimercaprol
(C) Calcium folinate
 (B) D-Penicillamine
(D) EDTA

116. Ferrous fumarate is assayed by
(A) Acidimetry
(B) Non-aqueous titration
(C) Cerimetry
(D) Precipitation titration
117. Sodium acetate is assayed by
(A) Precipitation titration
(B) Non-aqueous titration
(C) Gravimetry method
(D) Redox titration
118. Ringer's solution contains sodium chloride, potassium chloride and
(A) Magnesium chloride
(B) Calcium chloride
(C) Ferric chloride
(D) Zinc chloride
119. Ostwald's dilution law holds good only for —————
(A) weak electrolytes
(B) all electrolytes
(C) strong electrolytes
(D) strong acids and strong bases
120. The Van't Hoff equation for n moles of solute dissolve in V litres of solution is
(A) $\pi = nRT$
(B) $\pi V = nRT$
(C) $\pi P = nRT$
(D) $\pi P = nRT/V$
121. Insects can walk on the surface of water due to
(A) Viscosity
(B) Refractivity
(C) Surface tension.
(D) Optical activity
122. The external shape of the crystal is called
(A) amorphous
(B) habit
(C) plane of symmetry
(D) lattice

123. _____ reacts with water in perchloric acid and acetic acid, which makes the mixture anhydrous.
- (A) Acetic anhydride (B) Chloroform
(C) Acetone (D) Petroleum ether
124. According to Lowry Bronsted theory, acid is a
- (A) Proton donar (B) Proton acceptor
(C) Electro pain donar (D) Hydroxyl ion donar
125. Pfeffer's method is used to determine the
- (A) Surface tension (B) Density
(C) Vapour pressure (D) Osmotic pressure
126. Which of the following is Amphiprotic solvent used in Non-Aqueous titrations?
- (A) Pyridine (B) Perchloric acid
(C) Acetic acid (D) Toluene
127. Which of the following is a pM indicator?
- (A) Mordant black II (B) Crystal violet
(C) Starch iodide paper (D) Methyl orange
128. Which one of the following is the correct expression of Ostwald's dilution Law?
- (A) $K_c = \frac{\alpha^2}{(1-\alpha)V}$ (B) $K_c = \frac{\alpha^2 V}{1-\alpha}$
(C) $K_c = \frac{(1-\alpha)V}{\alpha^2}$ (D) $K_c = \frac{(1-\alpha)}{\alpha^2}$
129. Which one of the following is a mathematical expression of Raoult's law?
- (A) $\frac{P}{P-P_s} = \frac{n}{n+N}$ (B) $\frac{P-P_s}{P} = \frac{n}{n+N}$
(C) $(P-P_s)P = \frac{n}{n+N}$ (D) $\frac{P_s-P}{P} = \frac{n+N}{n}$

130. Optical activity is measured with the help of an instrument known as

- (A) Refractometer (B) Viscometer
 (C) Polarimeter (D) Conductometer

131. Which one of the following compounds will be optically active?

- (A) Succinic acid (B) Meso tartaric acid
 (C) Lactic acid (D) Chloro acetic acid

132. How many optical isomers are possible for Lactic acid?

- (A) 2 (B) 4
(C) 6 (D) 8

133. Geometrical isomerism is shown by

- (A) Lactic acid (B) Maleic acid
(C) 1-Butene (D) 1,1-Dichloro ethylene

134. Relative configuration of a chiral molecule is based on

- (A) Structure of D-Glucose
(B) Structure of D-Glycerol
 (C) Structure of D-Glyceraldehyde
(D) Structure of D-Fructose

135. Alkenes show geometrical isomerism due to

- (A) Asymmetry
(B) Resonance
(C) Rotation around single bond
 (D) Restricted rotation around a double bond

136. Geometrical isomers differ in _____ their physical and in _____ of their chemical properties.
- (A) all and many (B) all and all
(C) many and many (D) many and all
137. The solution which do not obey Raoult's Law are called
- (A) Ideal solution
 (B) Non ideal solution
(C) Buffer solution
(D) Non buffer solution
138. The compounds which have positive enthalpies of formation are called as
- (A) Enthalpy of formation
(B) Exothermic compound
 (C) Endothermic compound
(D) Standard enthalpy of formation
139. Reaction which are accompanied by the absorption of heat are called
- (A) Exothermic reaction (B) Endothermic reaction
(C) Heat constant (D) Constant volume
140. The Equation which indicates the amount of heat (enthalpy) changes in the reaction or process is called as.
- (A) Thermochemical equation
(B) Kirchoff equation
(C) Gibb's equation
(D) Hess equation
141. Co-factor for transamination is
- (A) Thymine (B) Riboflavin
 (C) Pyridoxine (D) Niacin

142. Provitamin D₂ is known as

- (A) Ergosterol
- (B) 7 - dehydro cholesterol
- (C) Ergo calciferol
- (D) Chole calciferol

143. In Knorr synthesis of pyrrole, _____ is used as a reactant.

- (A) Ethyl Aceto acetate
- (B) Diethyl malonate
- (C) Diethyl amine
- (D) Triethyl malonate

144. Aldehydes (Or) Ketones React with _____ to give Alcohols.

- (A) EAA (Ethyl Aceto Acetate)
- (B) DEM (Diethyl Malonate)
- (C) GR (Grignard Reagent)
- (D) TEA (Triethyl Amine)

145. Ortho and para - nitro toluenes are best synthesised and separated as a pure product by use of

- (A) Grignard reagent
- (B) Diazonium salts
- (C) Aceto acetic esters
- (D) Volhard reagents

146. Which of the following is a Grignard reagent?

- (A) Hydrogen bromide
- (B) Calcium bromide
- (C) Methyl magnesium bromide
- (D) Magnesium bromide

147. The energies of the orbitals increases in the following order
- (A) $1s < 2s < 3s < 3p < 4s < 3d < 4p$
 - (B) $1s < 2s < 3s < 3p < 3d < 4s < 4p$
 - (C) $4p < 4s < 3d < 3p < 3s < 2s < 1s$
 - (D) $4p < 3d < 4s < 3p < 3s < 2s < 1s$
148. Mass number is equal to
- (A) Number of protons + Number of electrons
 - (B) Number of protons + Number of neutrons
 - (C) Number of protons
 - (D) Number of electrons
149. In S_N^1 reaction the rate of the reaction depends on
- (A) Substrate only
 - (B) Nucleophile only
 - (C) Substrate and nucleophile
 - (D) Directly on catalyst
150. In an E2 reaction involving an alkyl halide and a base, the rate of the E2 reaction
- (A) depends linearly on the concentration of both reactants
 - (B) is independent of the concentration of both
 - (C) depends linearly on the concentration of the alkyl halide only
 - (D) is independent of the concentration of the alkyl halide
151. An E2 reaction occurs in _____ step(s).
- (A) Two
 - (B) Single
 - (C) Three
 - (D) Four
152. All negatively charged species are
- (A) Electrophiles
 - (B) Lipophiles
 - (C) Nucleophiles
 - (D) Hydrophiles

153. Of all the alkylhalides, which one is the most reactive in an E1 reaction
(A) alkyl fluorides (B) ~~alkyl iodides~~
(C) alkyl bromides (D) alkyl chlorides
154. The topical antifungal (terbinafine) is active by virtue of their ability to
 (A) Block squalene epoxide synthesis leading to defective cell membrane
(B) Inhibit synthesis of 1, 3, β - glucan
(C) Inhibit cell mitosis
(D) Bind to fungal cell membrane to cell hemolysis
155. Ethionamide is a derivative of
(A) Iso nicotinic hydrazide (B) ~~Iso nicotin amide~~
(C) Iso ethambutol (D) Iso quinol
156. Benzimidazole anthelmintics binds selectively to _____ of nematodes, cestodes and fluke worms.
(A) α - tubulin (B) γ - tubulin
 (C) β - tubulin (D) δ - tubulin
157. Levamisole contains which basic nucleus in it?
(A) Imidazole only (B) Thiazole only
(C) Indole only (D) Imidazo thiazole
158. Which of the following is used in treatment of filariasis (Wuchereria Bancrofti)?
 (A) Di ethyl carbamazine citrate (B) Albendazole
(C) Thiabendazole (D) Prazi quantel
159. The drugs that act against parasites in humans are called as
(A) Anti fungals (B) Anti bacterials
(C) Anti emetics (D) Anthelmintics

160. The drug Dipronhydramine hydrochloride is
(A) Imidazoline derivatives
(B) Thiophene derivatives
(C) Ethylene diamines derivative
(D) Aminoalkyl ethers derivative
161. Membrane stabilizing agents otherwise called as
(A) Calcium channel blockers
(B) Depolarization prolongators
(C) β - adrenergic blockers
(D) Sodium channel blockers
162. Which of the following drugs is an antimalarial?
(A) 9-aminoacridine
(B) Tolazamide
(C) Glibenclamide
(D) Acetohexamide
163. Which one is second generation non sedating Antihistamines
(A) Phenindamine tastrate
(B) Terfenadine
(C) Triprolidine hydrochloride
(D) Chlorpheniramine maleate
164. IUPAC Name for Histamine
(A) 2 - Dimethylamino ethyl Imidazole
(B) α - methyl benzyl Imidazole
(C) 4 - (2-aminoethyl) Imidazole
(D) 4 - (2-aminomethyl) Imidazole
165. Amlodopine a calcium channel blocker belongs to
(A) 1, 4 Dihydro pyridines
(B) 2, 4 Dihydro pyridines
(C) 3, 5 Dihydro pyridines
(D) 4, 6 Dihydro pyridines

166. Which of the following Antimalarial drug belong to 9-amino acridines?
- (A) Mepacrine
(B) Quinine
(C) Primaquine
(D) Amodia quine
167. Metoprolol, a selective β_1 - Adrenergic Blocker belong to _____ Anti hypertensives.
- (A) 1st generation
(B) 2nd generation
(C) 3rd generation
(D) 4th generation
168. Which of these compounds are unsaturated seven membered rings containing one nitrogen atom?
- (A) Azepine
(B) Pyridine
(C) Lanthine
(D) Furan
169. Most of the antimalarials possess _____ nucleus
- (A) Quinoline
(B) Thiazine
(C) Pyridine
(D) Piperazine
170. When Estrone is distilled with Zn dust, it forms
- (A) Chrysene
(B) Estrodiol
(C) Octa hydrocortisene
(D) Progesterone
171. A solution of cholesterol in chloroform, when treated with concentrated sulphuric acid develops a _____ colour in the chloroform layer
- (A) Blue
(B) Green
(C) Red
(D) Violet

172. _____ refers to the critical formation and storage of glycogen from glucose in the liver and muscle.
- (A) Glyconeogenesis
(B) Glycogenesis
(C) Glycolysis
(D) TCA cycle
173. Which one of the following is a female sex hormone?
- (A) Androstenediol
(B) Androstenolone
(C) Oestrogen
(D) Androgens
174. Pregnanediol on oxidation followed by reaction with Bromine and dehydrobromination yields
- (A) Pregnane
(B) Progesterone
(C) Ethisterone
(D) Androgen
175. Which of the following is a biologically active peptide?
- (A) Collagen
(B) Keratins
(C) Bradykinin
(D) Elastin
176. Which one of the following is a pyridine alkaloid?
- (A) Connine
(B) Hygrine
(C) Reserpine
(D) Quinine
177. Ephedrine undergoes oxidation with strong oxidising agents to give
- (A) Salicylic acid
(B) Acetyl Salicylic acid
(C) Benzoic acid
(D) Cinnauric acid
178. Fructose on treatment with phenylhydrazine and sodium acetate yields
- (A) Oza zone Sorbic acid
(B) Sucrose
(C) Fructosazone
(D) Fructo acetate

179. In spectro fluorimetry, absorption of emitted radiation by other analyte molecules called
- (A) Quenching
 - (B) Inner – filter effect
 - (C) Tyndall effect
 - (D) Raman scatter
180. Development time is shorter in
- (A) Paper chromatography
 - (B) Thin – layer chromatography
 - (C) High performance thin layer chromatography
 - (D) Two dimensional paper chromatography
181. In electromagnetic radiation, colorimetric (visible) range is from _____ to _____ nm.
- (A) 190 to 360
 - (C) 370 to 750
 - (B) 800 to 1200
 - (D) 2000 to 2400
182. In UV spectrophotometry, first derivative spectrum is a plot of the rate of change of absorbance with wave length against
- (A) Absorptivity
 - (B) Concentration
 - (C) Wave length
 - (D) Absorbance
183. During chemical derivatisation of amine, it is first diazotised with an aqueous solution of
- (A) Nitric acid
 - (B) Nitrous acid
 - (C) Nitrous oxide
 - (D) Sodium nitrite

184. A shift of λ_{\max} to longer wavelength is called as
- (A) Blue or hypsochromic shift
 - (B) Red or Bathochromic shift
 - (C) Hyper chromic shift
 - (D) Hypo chromic shift
185. The solvent kept in the tank moves up the thin layer of the solid adsorbent on the plate due to
- (A) Absorption
 - (B) Gravity
 - (C) Capillary action
 - (D) Reverse phase
186. Proteins and nucleotides can be separated by using cross linked
- (A) Cellulose
 - (B) Dextran gels
 - (C) Calcium silicate
 - (D) Polyamide
187. In column chromatography, the individual components of a mixture are separated by eluting the column with fresh solvent is
- (A) Elution analysis
 - (B) Frontal analysis
 - (C) Displacement analysis
 - (D) Reversible analysis

188. Photometric detector used in HPLC is

- (A) Diode – array detector
- (B) Refractive index detector
- (C) Electrochemical detector
- (D) Fluorescence detector

189. In IR, C = O stretching absorbs at

- (A) 1000 cm^{-1}
- (B) 1700 cm^{-1}
- (C) 1200 cm^{-1}
- (D) 1100 cm^{-1}

190. In IR, C = C and C \equiv N stretching vibration are located in the range of

- (A) $1500 - 1000\text{ cm}^{-1}$
- (B) $1000 - 900\text{ cm}^{-1}$
- (C) $1690 - 1600\text{ cm}^{-1}$
- (D) $1500 - 700\text{ cm}^{-1}$

191. The IR spectra is due to

- (A) Vibrational transitions
- (B) Electronic transitions
- (C) Rotational transitions
- (D) Spin reversal of Nuclei

192. In the titration of strong acids and strong bases by conductometric method, there is always an initial _____ in conductance
- (A) Increase
(B) Decrease
(C) Constant
(D) Non-linear
193. While performing radio immuno assay, the lowest concentration of a compound which can be detected in undiluted body fluid is called
- (A) Quantitation limit
(B) Cut-off level
(C) Spin-off level
(D) Titre
194. The most vital equipments essentially required for Radioimmuno Assay (RIA) are
- (A) Emulsifier and radioactive counters
(B) Centrifuge and separator
(C) Centrifuge and ball mill
(D) Centrifuge and radioactive counters
195. The electron bombardment in mass spectroscopy with energy 10 – 15 eV usually removes _____ electron(s) from the molecule
- (A) Three
(B) One
(C) Four
(D) Five

196. Chemical shift is expressed in one of the following units of

- (A) PPM (B) Ml
(C) RPM (D) cm^{-1}

197. ^{12}C nucleus in NMR spectroscopy is

- (A) Active
(B) Magnetically active
(C) Not active
(D) Not magnetically active

198. Faraday cup collector is employed in

- (A) IR
(B) NMR
(C) Mass
(D) UV

199. Using nitrogen rule, predict whether a molecule having mass number of 45 has

- (A) 4 Nitrogen (B) 2 Nitrogen
(C) 1 Nitrogen (D) No Nitrogen

200. The number of unsaturated centres in the molecule is calculated by

- (A) Double focussing
(B) Index of hydrogen deficiency
(C) Inductive effect
(D) Simple cleavage



PHARMACY

INDIA

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK



PHARMACY

INDIA

SPACE FOR ROUGH WORK

10/10/18



PHARMACY

INDIA

SEAL